

COMMENTARY

SPECIAL ISSUE

SENSORY PERCEPTION IN A CHANGING WORLD

Chemical cues and molecular mechanisms suspected in abiotic stress communication

Jiao Li¹, Lauric Feugere², Joerg Hardege³, Sofia Vámos¹ and Katharina C. Wollenberg Valero^{1,4,*}

ABSTRACT

For nearly a century, scientists have tried to resolve the sensory physiology of chemical communication caused by predation stress. Only recently have we evidenced that abiotic stressors from a changing world, such as heat and ocean acidification, also trigger chemical communication between aquatic organisms – which we dubbed abiotic stress communication. Generally, the behavioural and physiological response to stress-induced cues are well understood, whereas the molecular mechanisms – cue identities, pathways of release, and perception – of this stress communication remain unresolved. Here, we propose a framework to organize the existing evidence for candidate mechanisms involved in abiotic stress-induced chemical communication, focusing on heat and acidification as two major abiotic stressors with environmental relevance. Drawing on transcriptomic, metabolomic and behavioural evidence, we propose that stressor-specific communication likely involves multiple cues and parallel routes rather than a single mechanism, such as membrane-related processes. We call for integrative work that links -omics with chemical profiling and ecological function assays to uncover the mechanisms of abiotic stress communication.

KEY WORDS: Transcriptomics, Metabolomics, Sensory biology, Chemical communication

Introduction

Inter-individual communication is central to life, from bacteria (Taga and Bassler, 2003; Federle and Bassler, 2003) to plants (Gagliano et al., 2012) and animals (Laidre and Johnstone, 2013). One prominent form, chemical communication, involves the release of molecular cues that modify receiver behaviour or physiology (Regnier, 1971; Amo and Bonadonna, 2018; Sbarbati and Osculati, 2006). These cues mediate processes from predator avoidance to reproduction and social organization (Ohnstad et al., 2024; Roggatz et al., 2022; Fletcher et al., 2022; Kamio et al., 2022; Burks and Lodge, 2002). Chemical communication involves the production of a chemical cue, its release and transport into the environment, and its reception and integration by a receiver to trigger a response (Roggatz et al., 2022; Wyatt, 2014).

Chemical communication occurs in all environments, but is especially important in freshwater and marine aquatic environments where the reliance on visual cues is limited (Hay, 2009; Roggatz et al., 2022).

Stress communication describes how stressed organisms release cues affecting others (Abreu et al., 2016; Barcellos et al., 2011; Feugere et al., 2021b). Most of the knowledge on stress chemical communication comes from research on alarm systems caused by biotic stressors, such as predation and injury (Crane et al., 2022). On land, insects emit alarm pheromones eliciting antipredator behaviours (Verheggen et al., 2010), and similar mechanisms occur in birds, rodents and humans (Brandl et al., 2022; Carnevali et al., 2020; Bombail, 2019; Bombail et al., 2018). Fish can recognize conspecifics by smell (Solomon, 1977, citing Wrede, 1932), and injured fish release alarm cues provoking defensive behaviours in receivers (Brown and Smith, 1997; Chivers and Smith, 1998). This was first described as Schreckstoff in minnows (von Frisch, 1938, 1942). Alarm systems have since been documented in arthropods (Verheggen et al., 2010), gastropods (Atema and Stenzler, 1977), echinoderms (Snyder and Snyder, 1970), amphibians (Hews, 1988) and fishes (Brown, 2003; Wisenden et al., 2004). Even non-injured animals emit disturbance cues upon being startled (Crane et al., 2022; Jordão and Volpato, 2000; Ferrari et al., 2010). Known alarm or disturbance cues include hypoxanthine-3-N-oxide (Brown et al., 2004), oxysterol sulphate and 5 α -cyprinol sulphate (Li et al., 2024), and the glycosaminoglycans (GAGs) chondroitin sulphate (Mathuru et al., 2012) and keratan sulphate (Souza et al., 2007). Chemical communication is mostly studied in biotic stress contexts. Conversely, far less is known about whether abiotic stressors, such as warming and acidification, can likewise trigger inter-individual chemical stress communication. Despite decades of research since the first discoveries of biotic (von Frisch, 1938) stress chemical communication, the molecular identity and mechanisms of stress-induced chemical cue bouquets remain largely unknown (Stensmyr and Maderspacher, 2012). It also remains unclear whether abiotic and biotic stress cues and their pathways overlap. This represents a significant knowledge gap in the field of chemosensory biology and raises the important question of how animals might collectively respond to rapid global changes if chemical communication modulates species stress responses.

This knowledge gap, and two prior studies that observed behavioural changes in zebrafish and crayfish (Hazlett, 1985; Abreu et al., 2016), led us to the recent discovery that chemical communication extends to abiotic stressors. We showed that heat- or pH-stressed aquatic animals release cues that alter the growth, behaviour and gene expression of their receivers. This ripple effect propagates and amplifies stress responses within animal groups (Feugere et al., 2021a,b, 2023, 2025a). Prior to this, abiotic-related stress communication had only been anecdotally reported through behavioural studies in heat-stressed crayfish (Hazlett, 1985) and pH-stressed zebrafish (Abreu et al., 2016). In the context of rapid

¹OdysysLab, University College Dublin, School of Biology and Environmental Science, Belfield Campus, Dublin 4, Ireland. ²Marine Ecological and Evolutionary Physiology (MEEP) laboratory, Département de Biologie, Chimie et Géographie, Université du Québec à Rimouski, 300 Allée des Ursulines, Rimouski, QC G5L 3A1, Canada. ³School of Natural Science, University of Hull, Cottingham Road, Kingston-upon-Hull HU67RX, UK. ⁴Conway Institute, University College Dublin, Belfield Campus, Dublin 4, Ireland.

*Author for correspondence (katharina.wollenbergvalero@ucd.ie)

© J.L., 0000-0002-5716-5514; L.F., 0000-0002-4984-9350; J.H., 0000-0002-7815-3969; S.V., 0009-0001-9354-9603; K.C.W.V., 0000-0001-8858-1804

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

climate change, species will be increasingly exposed to abiotic stressors (Halpern et al., 2015; IPCC, 2019), especially higher temperatures owing to ocean warming and lower pH levels due to ocean acidification (Roggatz et al., 2022). Ocean acidification is caused by rising atmospheric CO₂ released from human activities, which lowers seawater pH globally (Doney et al., 2009). Ocean acidification and ocean warming do not only manifest by average shifts, but also by increasing diel pH and temperature fluctuations (Bednaršek et al., 2022). Consequently, intense events such as short-term acidification, marine heatwaves and their co-occurrence (Landschützer et al., 2018; Burger et al., 2022) are becoming increasingly frequent and are superimposed on gradual global changes (Landschützer et al., 2018; Burger et al., 2022). These time scales matter for the individual organism, which will experience acute stress during its life course, and especially early development, potentially triggering stress communication.

The developmental stage of organisms may fundamentally shape these pathways. Early life stages are capable of chemical communication (Lucon-Xiccato et al., 2020; Atherton and McCormick, 2015; Cao and Li, 2020). Embryos lacking a differentiated epidermis are more permeable, and may release chemical cues (but also uptake them) through loosely organized outer cell layers (Le Guellec et al., 2004). Moreover, the presence of a chorion limits the permeability for certain size classes of molecules (Pelka et al., 2017). Once the functional epidermis is developed, permeability to outgoing signals may be more limited or utilize other pathways, such as urine excretion.

In this Commentary, we discuss the evidence of abiotic stress communication from our previous studies on zebrafish embryos

(*Danio rerio*), sea bream (*Sparus aurata*) and marine ragworms (*Hediste diversicolor*; Feugere et al., 2021b, 2023; Beine et al., 2024) exposed to elevated temperature and acidified pH. We discuss possible molecular mechanisms and propose an evolutionary *raison d'être* for abiotic stress communication and its implication in a fast-changing world (Fig. 1). We have identified five mechanisms with evidence of involvement in abiotic stress chemical communication: (1) lipids and membranes, (2) proteins, (3) extracellular matrix (ECM), (4) other small molecules and (5) immune system entanglement (Table 1). We present these mechanisms in the next sections.

Lipids and membranes

Temperature-induced membrane remodelling as possible lipid cue release mechanism

Membrane lipid composition varies in response to environmental stressors through both homeoviscous and homeophasic adaptation maintaining their fluidity and phase stability (Jin et al., 2021; Sinensky, 1974; Vigh et al., 1998; Hazel, 1995, 1997). Warming lowers acyl chain unsaturation (Holm et al., 2022), and can cause shifts in lipid phase behaviour, including transitions to non-lamellar phases such as the inverse hexagonal phase. Heat stress can also alter headgroup composition and cholesterol content (Vigh et al., 1998; Hazel, 1997). For example, in *Tetrahymena*, heat increases membrane phosphatidylcholines (Nozawa, 2011), a lipid class derived from phosphocholine (Li and Vance, 2008). Likewise, experimental ocean warming and acidification treatments altered the muscle membrane lipidome of *Pandalus borealis* shrimp (Feugere et al., 2025b).

Therefore, we propose that lipid monomers deriving from stress-induced membrane remodelling could act as abiotic stress cues.

Increased temperature and acidification




Fig. 1. Schematic overview of the main molecular mechanism of abiotic stress communication.

Upon exposure to heat and acidification, pathways are activated in organisms that initiate the release of stress cues into the environment by senders. These are then perceived by receiver organisms, in which specific molecular pathways are triggered which alter properties of the receiver. This ripple effect propagates and amplifies stress responses between individuals. Created in BioRender. Wollenberg Valero, K. (2025) <https://BioRender.com/g635dq1>. This figure was sublicensed under CC-BY 4.0 terms.

Table 1. Five proposed molecular mechanisms of abiotic stress chemical communication

Molecular pathways	Hypotheses and evidence for their involvement in abiotic stress chemical communication
Lipids and membranes	
Release	Hypothesis: abiotic stressors remodel membranes, releasing lipids or membrane fragments. Evidence: <i>Hydra</i> uses extracellular vesicles (ECVs) for social signalling through the medium.
Cues	Hypothesis: abiotic stress cues are lipids released under abiotic stress. Evidence: lipids are known cues in chemical communication. Medium from heat-stressed zebrafish embryos contains phosphocholines and phosphatidylcholines.
Reception	Hypothesis: membrane-bound lipids sense stress cues. Evidence: receivers of heat-stress cues in zebrafish upregulate <i>prg4a</i> (linked to phosphocholine) and the scramblase <i>xkr8.2</i> (which flips membrane phospholipids). <i>xkr8.2</i> is upregulated in lateral line hair cells, which may detect chemical cues. Ragworms receiving cues from pH-stressed fish differentially express <i>gpr45</i> , a lipid-signaling G-protein coupled receptor (GPCR).
Proteins	
Release	Hypothesis: proteins such as ABC transporters secrete abiotic stress cues. Evidence: proteins including ABC transporters export signalling cues across kingdoms and are involved in the cellular stress response. No evidence yet for their role in abiotic stressor cue release.
Cues	No evidence for the involvement of proteins/peptides in abiotic stress signalling.
Reception	Hypothesis: chemosensory, transport and receptor proteins are involved in abiotic stress cue reception. Evidence: zebrafish embryos exposed to heat stress and to heat-stressed conspecific cues upregulated chemosensory receptor genes (<i>ora3</i> , <i>olfcb1</i>). Marine ragworms exposed to pH-stressed fish cues modulated transcription of several receptors (<i>trpc6</i> , <i>otop</i> , <i>otog</i> , <i>gbra6</i> and several GPCRs incl. <i>gpr45</i> , <i>fmar</i>), transport (<i>slc4a10</i>) and ABC transporter (<i>abcb6</i>) proteins.
Extracellular matrix (ECM)	
Release	Hypothesis: ECM secretes abiotic stress cues. Evidence: heat stress alters the expression of ECM-related genes linked to glycosaminoglycans (GAGs). GAGs are alarm cues in fish.
Cues	Hypothesis: abiotic stress cues are ECM components. Evidence: ECM components such as hypoxanthine-3-N-sulphate and chondroitin sulphate are alarm cues. There is no evidence for their roles in abiotic stress signalling, suggesting that biotic and abiotic stress communication may differ.
Reception	Hypothesis: ECM detects abiotic stress cues. Evidence: zebrafish embryos exposed to abiotic stress cues upregulated an ECM adaptor linking proteoglycans (<i>matrilin 3a</i>), and genes linked to GAG metabolism (<i>krt17</i> , <i>krt97</i> , <i>chs1</i> , <i>chst2b</i> and <i>prg4a</i>).
Other small molecules	
Release	Hypothesis: small molecules pass through membranes and lead to abiotic stress cues. Evidence: heat-stressed zebrafish embryos activate small molecule metabolism including pentose phosphate (PPP) and methionine metabolism pathways (involving <i>gpia</i> and <i>pgm1</i>).
Cues	Hypothesis: abiotic stress cues are small molecules. Evidence: the medium from heat-stressed zebrafish embryos contain primary metabolites, including carbohydrates (PPP), sulphur-containing compounds (transsulphuration pathway; e.g. 3-mercaptolactate).
Reception	No evidence for the role of specific small molecules in receiving abiotic stress cues.
Immune system	
Release	No evidence for immune system involvement in abiotic stress cue release.
Cues	Hypothesis: abiotic stress cues stem from immune responses. Evidence: damage-associated molecular patterns (DAMPs; e.g. oxidised lipids or degraded ECM fragments) can mimic pathogen-associated molecular patterns (PAMPs) and activate innate immunity via epithelial detection and inflammatory signaling. DAMPs are often purine derivatives. The alarm cue hypoxanthine (Schreckstoff), a purine derivative, may activate immune responses in receivers.
Reception	Hypothesis: abiotic stress cues induce immune pathways. Evidence: zebrafish embryos exposed to heat stress cues downregulated genes involved in immunity signalling (<i>si:ch211-214b16.2</i> and <i>il-1β</i>).

We propose the mechanisms underlying release of the abiotic stress cues by the sender, their molecular identity and, lastly, reception by the receiver. Detailed references for each point are given in the main text.

However, their amphiphilic nature limits their persistence in water, and they tend to form micelles or surface monolayers (Pichot et al., 2013). A different mechanism of transport in aquatic environments would require monomers to be stabilized by carrier proteins (Mangiacotti et al., 2023), or to self-assemble into micelles or vesicles (spheres of lipid bilayers, extracellular vesicles or ECVs). ECVs may carry and release other chemical cues as cargo into the environment. The density of nano-sized ECVs is around 1.10–1.19 g ml⁻¹ (van der Vlist et al., 2012), compared with 1.03 g ml⁻¹ for seawater (depending on temperature and/or salinity; McDougall and Parker, 2011) and 1 g ml⁻¹ for freshwater. Therefore, ECVs should remain suspended in seawater but slightly sink in freshwater, making them likely candidates to transport signalling molecule cargo between organisms in both media. In fact, the aquatic invertebrate *Hydra vulgaris* has been shown to release ECVs

containing molecular messages for other individuals into its culture medium (Moros et al., 2021).

Lipids and derivatives as abiotic stress cues

Zebrafish embryos kept for 24 h post-fertilization (hpf) under a fluctuating heat stress activated two key pathways (glycerophospholipid and ether lipid metabolism) and released lipids into the medium. These included fatty acyls, glycerolipids, sphingolipids, glycerophospholipids and steroid esters (Feugere et al., 2023), which links membrane lipid turnover to heat stress signalling. Among these lipids, phosphocholine and glycerophosphocholine (a derivative of phosphocholine) stood out. Both of these lipids were previously shown to be upregulated in heat-stressed fish tissues (Karakach et al., 2009; Zhang et al., 2022). Lipids, such as sphingolipids, are already known for protecting

cells under heat stress and acting as molecular messengers (Balogh et al., 2013; Török et al., 2014), mediating mating in reptiles (Goldberg et al., 2017), guiding larval migration in lampreys (Li et al., 2018), communicating the presence of copepod predators to dinoflagellate prey (Selander et al., 2015), and marking group identity in catfish (Matsumura et al., 2007). Furthermore, lipids can serve as coral settlement cues (Jorissen et al., 2021). In zebrafish, the yolk sac likely serves as an abundant source of such lipids (Miyares et al., 2014). Moreover, crustacean pheromones and contact cues may be simple, heat-stable lipids (Takahashi, 1974; Zhang et al., 2011). Altogether, there is strong evidence that abiotic stress can turn membrane lipids into messages.

Uptake of lipids from abiotic stress medium

Receptors for free phospholipids in aquatic environments are largely unknown. Although lipid head group exposure can trigger cell-to-cell signals (e.g. in apoptosis), this mechanism is understood intracellularly (Fadok et al., 1998), but under-researched in inter-organismal communication. However, lipid signals can integrate into cell membranes through membrane fusion and alter fluidity, curvature or receptor distribution, making membranes themselves interpreters of environmental cues (Vigh et al., 1998). Contact with ECVs could bulk-transfer metabolic and stress-related information. In zebrafish, sensory cells such as those in the lateral line might detect environmental changes via lipid uptake. For instance, 24-hpf zebrafish embryos exposed to thermal stress cues upregulated genes involved in phospholipid and anion transport. This included the scramblase *xkr8.2*, which flips membrane phospholipids and is both highly expressed in lateral line hair cells and upregulated by stress cues (Feugere et al., 2023; Elkon et al., 2015). Moreover, multiomics analyses functionally linked a gene associated with the ECM (see below), *prg4a*, to phosphocholine, and its consistent upregulation across several independent experiments strengthens its candidate role in cue response (Feugere et al., 2023). In marine ragworms grown in a future ocean regime (adjusted diel and seasonal temperature and pH cycles), *gpr45*, a lysophosphatidic acid-like lipid-signalling G-protein coupled receptor (GPCR), is differentially expressed in response to exposure to the metabolites of *Sparus aurata* sea bream exposed for an hour to acidified pH (Marchese et al., 1999; Cui et al., 2016; Geraldo et al., 2021; Kiss et al., 2012; Lee et al., 2023; Beine et al., 2024). This shows that lipid signals present in abiotic stress media can be taken up by receivers, again supporting their role in abiotic stress communication.

Proteins

Protein-mediated abiotic stress cue release

Although there is no evidence for secretory proteins actively transporting abiotic signalling cues into the aqueous environment, some protein candidates warrant further consideration. For example, ATP binding cassette (ABC) transporters are a ubiquitous large family of proteins that use ATP to actively transport molecules across membranes (Theodoulou and Kerr, 2015), including ions, organic molecules of different polarity, lipids, peptides and small proteins. Beyond individual molecule export, some ABC subfamilies are also implicated in vesicle formation or cargo loading, suggesting they might facilitate cue packaging into lipid vesicles or microdomains (Theodoulou and Kerr, 2015). ABC transporters are known to export signalling cues across kingdoms. In tunicates, they guide sperm cell motility by exporting signalling lipids (Kassmer et al., 2020). In plants, they mediate phytohormone transport within tissues and into the soil (Borghi et al., 2015; Sasse et al., 2015), support chemical kin recognition (Biedrzycki et al., 2011; Bailly, 2014) and facilitate the

release of small volatile organic compounds (Adebisin et al., 2017). ABC transporters also contribute to the heat stress response. In yeast, heat-induced ABC transporters deliver ergosterol to plasma membranes, increasing their fluidity and integrity (Godinho et al., 2021). In pigeon pea, ABC genes are directed by stress-responsive promoters, to export bioactive compounds such as hormones and signalling molecules (Niu et al., 2021).

Protein-mediated stress cue uptake

GPCRs, vanilloid receptors [transient receptor potential (TRP) channels], scavenger receptors and GABA receptors mediate pheromone and kairomone detection (Fortes-Marco et al., 2013), environmental sensing and systemic physiology regulation, making them strong candidates for stress cue detection. In adult fish, the olfactory bulb processes chemical cues (Hamdani and Døving, 2007), but this system is still maturing in embryos and larvae (Li et al., 2005). Zebrafish express three main olfactory GPCRs: olfactory receptors (Liberles and Buck, 2006; Hashiguchi and Nishida, 2007; Liberles, 2009), trace amine-associated receptors, vomeronasal receptors (V1Rs, V2Rs), plus other pheromone-detecting GPCRs (Pfister and Rodriguez, 2005). Cues binding to GPCRs activate G-protein pathways, and open ion channels via increased cAMP (Kandel et al., 2000). The lateral line system expresses TRP and purinergic chemoreceptors (Corey et al., 2004; Xiang et al., 1999; Quick et al., 2012; Muñoz et al., 1999), and also detects chemical cues via mechanosensory hair cells (Coombs et al., 2013; Dijkgraaf, 1963; Webb, 2013). Recently, 45 novel chemoreceptors were discovered in zebrafish hair cells, including 19 olfactory GPCRs, TRPs and pheromone receptors (Desban et al., 2022). We also found that zebrafish embryos exposed to heat and heat-induced stress cues simultaneously, but not to heat alone, altered the expression of the chemosensory receptor genes *ora3* and *olfcb1* (Feugere et al., 2023). In marine ragworms, pH-stress-exposed fish cues upregulated TRPC6, downregulated OTOG and OTOG, and modulated GBRA6 and several GPCRs such as GPR45 and FMAR (Beine et al., 2024; Bernal-Ero et al., 2014). These patterns suggest responses to stress cues induced by altered extracellular pH or heat stress, either directly or indirectly. In marine ragworms, exposure to stress cues from pH-stressed sea bream induced distinct gene expression patterns (Beine et al., 2024). For instance, stress cues upregulated *slc4a10* whereas direct pH stress downregulated it. Notably, stress cues downregulated *abcb6* (Ye et al., 2020; Rakvács et al., 2019), an ABC transporter with membrane phospholipid binding function (Lee et al., 2023). Owing to their secretory nature, ABC transporters are unlikely to be receptors for abiotic stress cues, but they could function in intracellular reprogramming following the detection of stress cues. However, ABC transporters can internalize quorum sensing signals in microorganisms (Taga and Bassler, 2003), or even 'steal' xenosiderophores, which are iron-binding molecules secreted by other species (Liu and Kakeya, 2020). Besides their signalling role being mostly known from plants and microbes, the upregulation of ABC transporters in heat-stressed aquatic organisms may simply reflect a general cellular protection response (Lauritano and Ianora, 2020).

Extracellular matrix

Damage-induced release of stress cues

Alarm cues are associated with injury. Schreckstoff was first identified in skin extract triggering startle responses in conspecifics (Smith, 1979). von Frisch (1942) noted that Schreckstoff was only produced by epidermal cells. In fish, the ECM is essential for epithelial integrity and supports the mucus layer enriched in GAGs (Syed et al., 2022; Frantz et al., 2010). This supports the idea that

membrane damage caused by abiotic stress could cause the passive leakage of cytoplasmic components or unstructured membrane fragments, including parts of the ECM. This structure is tightly linked to cell membranes and includes proteoglycans known to carry potential cue components, including GAGs. Known ECM alarm cues include GAGs such as chondroitin sulphate and mucin proteins (Mathuru et al., 2012). The original Schreckstoff, hypoxanthine-3-N-oxide (Parra et al., 2009), is a chemical relative of hypoxanthine and a derivative of adenine. The latter is part of the ECM structure in zebrafish (Deis et al., 2025). Altogether, this supports the hypothesis that abiotic stressors lead to the release of GAGs via disrupted membrane fragments with the ECM attached. However, there was no evidence for specific ECM fragments involved in abiotic stress communication in our prior study in zebrafish (Feugere et al., 2023).

Extracellular matrix as abiotic stress cue receiver?

The ECM is not purely structural. It also modulates signalling through ligand binding and receptor presentation (Kim et al., 2011), making it a likely interface for receiving and releasing stress cues. In 24-hpf zebrafish embryos exposed to medium from heat-stressed conspecifics, we showed that heat stress and stress cues alter the expression of ECM-related genes, such as matrilin 3a (*matn3a*, an ECM adaptor linking proteoglycans; Feugere et al., 2023; Paulsson and Wagener, 2018), proteoglycan 4a (*prg4a*), and genes related to the GAGs keratan sulphate and chondroitin sulphate biosynthesis (*chs1*, *chst2b*, *krt17*, *krt97*) (Feugere et al., 2023). This suggests a role of ECM molecules in abiotic stress-mediated cue reception. Multiomics linked *prg4a* to phosphocholine, a lipid that is a potential heat stress cue, and was consistently upregulated across our experiments. In marine ragworm, receivers of stress cues from pH-stressed sea bream altered expression of four ECM-related genes (*ust*, *xylt2*, *b4galt7*, *b3galt4*), involved in chondroitin sulphate metabolism (Beine et al., 2024). Altogether, this supports the role of ECM in cue perception within and across different species.

A potential role for other small molecules

In addition, carbohydrates (and conjugates) were identified in the medium of heat-stressed zebrafish. These cues, such as deoxyribose, likely result from heat-induced metabolic shifts, including glycolysis and the key redox regulator pentose phosphate pathway (PPP). Genes associated with PPP (*gpi1* and *pgm1*) were upregulated in the heat-stressed zebrafish embryos (Feugere et al., 2023; Stincone et al., 2015; Karakach et al., 2009), which links a heat-induced metabolic process to its products identified in the stress medium. In fact, carbohydrates are known information carriers (Gabijs et al., 2004). Likewise, sulphur-containing compounds are known alarm cues (Brechtbühl et al., 2013; McLean et al., 2021; Miyazaki et al., 2006; Li et al., 2024) and two that are involved in methionine metabolism (Sörbo, 1987; Andreeßen et al., 2018) were identified in the thermal stress medium of zebrafish embryos (as 3-mercaptolactate and 2-oxo-4-methylthiobutanate; Feugere et al., 2023). 3-Mercaptolactate is synthesized by lactate dehydrogenase A (*ldha*), which we showed to be heat-induced (Feugere et al., 2023). 3-Mercaptolactate is excreted in urine and induces anxiety-like behaviours in mice (Nagahara et al., 2013; Akaoshi et al., 2020; Andreeßen et al., 2018; Song et al., 2019).

The question that follows is then, how can these small molecules be released from the thermally stressed individual? Metabolites, if they are hydrophobic, or polar and small, may readily diffuse through the cell membrane passively (Cooper, 2000). Two examples for passive diffusion of signalling molecules are the alkaloid neurotoxins

saxitoxin and tetrodotoxin. These molecules evolved as chemical defences, but also function as inter-organismal cues (Roggatz et al., 2019). Their ability to passively cross membranes allows them to act without specialized release pathways (Tikhonov and Zhorov, 2018; Van Theemsche et al., 2020). Similar to this model, smaller, more volatile metabolites are strong candidates for cues that can be detected at a distance from their source (Alberts and Werner, 1993; Konishi, 1970; Pozzer et al., 2022). In our previous experiment, classical alarm cues, including hypoxanthine-3-N-oxide, trigonelline and homarine (Poulin et al., 2018; Parra et al., 2009), were released by control zebrafish embryos rather than the heat-stressed ones (Feugere et al., 2023), suggesting chemical differences between abiotic stress and alarm cue bouquets. In summary, although the exact mechanism of small molecule cue release is unknown, and no evidence exists yet for the role of small molecules in abiotic stress communication, we showed that some are released into the medium by heat-stressed developing embryos as metabolic products of the primary metabolism.

Immune system entanglement

The immune system may misinterpret abiotic stress cues as pathogen-associated molecular patterns (PAMPs), triggering responses through shared pathways. Damage-associated molecular patterns (DAMPs), such as oxidized lipids or degraded ECM fragments, can mimic PAMPs, activating innate immunity via epithelial detection and inflammatory signalling (Tang et al., 2012; Varki, 2011). DAMPs are often purine derivatives (Russo and McGavern, 2015), such as hypoxanthine (Schreckstoff). In threespine stickleback, predator and alarm cues similarly modulated immune and signalling pathways (Sanogo et al., 2011). In zebrafish embryos, exposure to heat and heat stress cues downregulated *si:ch211-214b16.2*, likely a NOD2 ortholog (Feugere et al., 2023), and heat stress cues downregulated *il-1 β* involved in immune pathways and inflammation, possibly indicating immune suppression (Feugere et al., 2021b). Although NOD2 typically detects bacterial peptidoglycans, its modulation here may reflect recognition of stress-modified glycosaminoglycans in ECM-bound cues. Notably, *thr18*, a fish-specific toll-like receptor associated with lipid recognition and expressed in lateral line hair cells, was activated by stress cues of heat-exposed zebrafish embryos (Nie et al., 2018; Elkon et al., 2015; Feugere et al., 2023). Toll-like receptors might therefore recognize stress-induced lipids or ECM fragments, triggering MAPK or NF- κ B cascades and fear-like behaviours (Kirsten et al., 2018). Overall, this indicates that the reception of thermal stress cues modulates the immune system of receivers.

Why would abiotic stress communication exist?

In this Commentary, we describe the phenomenon of abiotic stress communication and discuss possible underlying molecular mechanisms. We have shown abiotic stress induces chemical communication, defined as information exchange, irrespective of intentionality, by means of chemical cues affecting their receivers (Wyatt, 2014). Zebrafish embryos receiving heat stress cues exhibited stress-like molecular responses, developed faster and were hypoactive, similar to heat-stressed conspecifics. Marine ragworms changed their burrowing, freezing and escaping behaviour when receiving cues from pH-stressed conspecifics as well as pH-stressed sea bream (Feugere et al., 2021a; Beine et al., 2024). The purpose of injury-induced biotic stress communication is likely for receivers to perceive the predation risk, without intentionality by the sender (Crane et al., 2022; Chivers et al., 2012). This would make them 'honest' biochemical byproducts of stress that unintentionally inform

others (Darwin, 1871; Andersson, 1994; de Groot and Smeets, 2017). Stress communication, whether biotic or abiotic, may in fact be widespread across social animal species owing to the evolutionary conservation of stress response mechanisms (Brandl et al., 2022). The evolutionary basis of abiotic stress communication is less obvious than for alarm signals caused by predation, where the threat is immediate and acute.

For abiotic stress communication, all mechanisms mentioned above similarly point at an unintentional and involuntary release of stress cues into the environment, originating from cell damage or changes in the primary metabolism of the sender. Therefore, we discuss four possible roles for abiotic stress communication, which are to: (i) be informed about others' physiological status, (ii) coordinate stress responses in a group and (iii) mitigate abiotic stress effects through either priming for the abiotic stressor or initiating stress responses, for instance by completing embryonic development faster or displaying specific behaviours (escaping, freezing, moving to another habitat). Lastly, (iv) abiotic stress cues may overlap with biotic stress cues and therefore cause similar reactions in receivers.

Role 1: physiological status

Akin to social stress transmission, abiotic stress communication may help to inform others of stress (Brandl et al., 2022). One may argue that there is no need to have a chemical stress communication system as the individual perceives the abiotic stress itself. Animals are known to gather as much information as possible, including socially acquired information, even if inadvertently (Dall et al., 2005). Species are also known to eavesdrop on alarm and disturbance cues of other species (Turner et al., 2023). Therefore, the ability for an animal to detect another animal's stress cues can inform it about its health status or whether a habitat is suitable.

Role 2: coordination of stress group responses

An abiotic stress communication system may be particularly useful to coordinate responses in social species (Grunst and Grunst, 2024). In a mechanism akin to quorum sensing (Li and Nair, 2012), a minimum threshold of stress cues may help the group coordinate its stress response to the threat, acting as honest social signals favouring group cohesion (Dall et al., 2005). This aligns with alarm and disturbance cue studies showing group responses depending on the stress experience of senders and receivers, with dose- and threshold-dependent patterns (Goldman et al., 2019, 2020). Likewise, as stress responses and stress perception vary among individuals (Guscelli et al., 2019; Melanson et al., 2023), abiotic stress communication may serve as a social coordination checkpoint that matches the group response to the level of the abiotic stressor, to coordinate responses such as developmental speed. This social buffering may counterbalance stress transmission to prevent maladaptive costs when environmental changes occur on longer time scales (discussed in Gilmour and Bard, 2022).

Role 3: mitigation of abiotic stress effects

Abiotic stress communication may be a hormetic mechanism, a low-intensity stressor priming the stress response and reducing the damage caused by additional abiotic stressors, such as improving DNA repair (Feugere et al., 2025a). Stress cues released by heat-stressed individuals accelerated growth in naive receivers and amplified it in heat-stress-exposed receivers (Feugere et al., 2021a, b, 2023). Abiotic stress communication may accelerate individual growth within cohorts to increase survival, as developed animals are more equipped to cope with the stress than developing ones

(Hamdoun and Epel, 2007; Cowan et al., 2024). Abiotic stress communication may prime animals through the activation of the molecular pathways, including the immune system defenses, needed to respond to the abiotic stressor.

Role 4: overlap with biotic stress cues

In our Commentary, we summarized multiple lines of evidence for membrane lipids and/or membrane fragments including the ECM being involved in abiotic stress communication. Similarly, lipids have multiple functions in social signalling, and the known injury-induced Schreckstoff components are most likely components of the ECM. Therefore, if abiotic stressor-induced membrane fragments come into contact with another organism, they may interpret the cue as alarm signals, as the same compounds are carried via the ECM or membrane lipids.

Future research directions

Abiotic stress communication appears across diverse organisms, tissues and developmental stages. Given that environments are increasingly challenging to organisms, it becomes crucial to further our understanding of its molecular mechanisms if abiotic stress communication has the potential to amplify stress responses. In zebrafish, early-life responses to stress cues may involve olfactory pathways, but confirmation in adults is still needed (Feugere et al., 2023). Mechanisms likely vary across taxa and stressors, e.g. heat, low pH and osmotic stress, yet converging patterns emerge. Chondroitin sulphate and other glycosaminoglycans recur across species, and many abiotic stressors affect membrane structure, implicating conserved pathways such as vesicle-mediated transport and ECM remodelling in cue propagation. Unlike alarm cues such as Schreckstoff, which are often species-specific (von Frisch, 1942), abiotic stress communication may rely on broadly shared components. This aligns with the concept of a cue bouquet composed of diverse molecules, including lipids, proteins, sugars or ECM fragments, acting in concert. In terrestrial species, lipids often require protein binding for signalling (Mangiacotti et al., 2023; Telser, 2002) and similar packaging into lipid vesicles enriched with small molecules and glycoproteins may analogously occur in aquatic systems. However, decades of research on Schreckstoff (e.g. Mathuru et al., 2012) have evidenced that it is extremely difficult to dissect the role of cues in signalling bouquets, calling for future research to identify what chemicals among stress metabolites act as effective stress cues driving abiotic stress communication. Acknowledging these limitations, we call for discussions among molecular biologists, ecophysiologicals, chemical ecologists and evolutionary scientists to resolve this outstanding question. A unified framework is needed to recognize chemical stress communication as a widespread, fundamental process. Uncovering more about this mechanism will require systemic, multiomic and behavioural approaches (Hebets et al., 2016; Brunetti et al., 2018).

Acknowledgements

We thank the organizers and participants of the Society of Experimental Biology symposium 'Sensory Perception in a Changing World – Impacts on Physiology and Behaviour' (Liverpool, 2025) for the interesting contributions and discussions.

Competing interests

The authors declare no competing or financial interests.

Funding

K.C.W.V., J.H. and L.F. acknowledge funding by the University of Hull toward the MolStressH2O doctoral research cluster. K.C.W.V., J.L., and S.V. acknowledge funding by the European Union (European Research Council, MolStressH2O, no. 101044202). Views and opinions expressed are however those of the author(s)

only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. Open Access funding provided by University College Dublin. Deposited in PMC for immediate release.

ECR Spotlight

This article has an associated ECR Spotlight interview with Jiao Li and Lauric Feugere.

Special Issue

This article is part of the special issue 'Sensory Perception in a Changing World', guest edited by Almut Kelber, Kathleen M. Gilmour and Sanjay Sane. See related articles at https://journals.biologists.com/jeb/issue/229/Suppl_1.

References

- Abreu, M. S., Giacomini, A. C. V., Gusso, D., Koakoski, G., Oliveira, T. A., Marqueze, A., Barreto, R. E. and Barcellos, L. J. G. (2016). Behavioral responses of zebrafish depend on the type of threatening chemical cues. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **202**, 895-901. doi:10.1007/s00359-016-1129-5
- Adebesin, F., Widhalm, J. R., Boachon, B., Lefèvre, F., Pierman, B., Lynch, J. H., Alam, I., Junqueira, B., Benke, R., Ray, S. et al. (2017). Emission of volatile organic compounds from petunia flowers is facilitated by an ABC transporter. *Science* **356**, 1386-1388. doi:10.1126/science.aan0826
- Akahoshi, N., Minakawa, T., Miyashita, M., Sugiyama, U., Saito, C., Takemoto, R., Honda, A., Kamichatani, W., Kamata, S., Anan, Y. et al. (2020). Increased urinary 3-mercaptoplactate excretion and enhanced passive systemic anaphylaxis in mice lacking mercaptopyruvate sulfurtransferase, a model of mercaptolactate-cysteine disulfiduria. *Int. J. Mol. Sci.* **21**, 818. doi:10.3390/ijms21030818
- Alberts, A. C. and Werner, D. I. (1993). Chemical recognition of unfamiliar conspecifics by green iguanas: functional significance of different signal components. *Anim. Behav.* **46**, 197-199. doi:10.1006/anbe.1993.1177
- Amo, L. and Bonadonna, F. (2018). Editorial: the importance of olfaction in intra- and interspecific communication. *Front. Ecol. Evol.* **6**, 71.
- Andersson, M. (1994). *Sexual Selection*. Princeton University Press.
- Andreeßen, C., Wolf, N., Cramer, B., Humpf, H.-U. and Steinbüchel, A. (2018). *In vitro* biosynthesis of 3-mercaptoplactate by lactate dehydrogenases. *Enzyme Microb. Technol.* **108**, 1-10. doi:10.1016/j.enzmictec.2017.08.005
- Atema, J. and Stenzler, D. (1977). Alarm substance of the marine mud snail, *Nassarius obsoletus*: biological characterization and possible evolution. *J. Chem. Ecol.* **3**, 173-187. doi:10.1007/BF00994144
- Atherton, J. A. and McCormick, M. I. (2015). Active in the sac: damselfish embryos use innate recognition of odours to learn predation risk before hatching. *Anim. Behav.* **103**, 1-6. doi:10.1016/j.anbehav.2015.01.033
- Bailey, A. (2014). Structure-function of plant ABC-transporters. In *Signaling and Communication in Plants* (ed. F. Baluška), pp. 219-240. Cham: Springer International Publishing.
- Balogh, G., Péter, M., Glaz, A., Gombos, I., Török, Z., Horváth, I., Harwood, J. L. and Vigh, L. (2013). Key role of lipids in heat stress management. *FEBS Lett.* **587**, 1970-1980. doi:10.1016/j.febslet.2013.05.016
- Barcellos, L. J. G., Volpato, G. L., Barreto, R. E., Coldebella, I. and Ferreira, D. (2011). Chemical communication of handling stress in fish. *Physiol. Behav.* **103**, 372-375. doi:10.1016/j.physbeh.2011.03.009
- Bednaršek, N., Beck, M. W., Pelletier, G., Applebaum, S. L., Feely, R. A., Butler, R., Byrne, M., Peabody, B., Davis, J. and Strus, J. (2022). Natural analogues in pH variability and predictability across the coastal Pacific estuaries: extrapolation of the increased oyster dissolution under increased pH amplitude and low predictability related to ocean acidification. *Environ. Sci. Technol.* **56**, 9015-9028. doi:10.1021/acs.est.2c00010
- Beine, K., Feugere, L., Fletcher, N., Power, M. L., Connell, L. J., Bates, A., Li, J., Winter, M. R., Sellers, G. S. and Mincarelli, L. F. (2024). The molecular arsenal of the key coastal bioturbator *Hediste diversicolor* faced with changing oceans. *bioRxiv*. doi:10.1101/2024.09.20.614147
- Berna-Erro, A., Albarran, L., Dionisio, N., Redondo, P. C., Alonso, N., Gomez, L. J., Salido, G. M. and Rosado, J. A. (2014). The canonical transient receptor potential 6 (TRPC6) channel is sensitive to extracellular pH in mouse platelets. *Blood Cells Mol. Dis.* **52**, 108-115. doi:10.1016/j.bcmd.2013.08.007
- Biedrzycki, M. L., V. L. and Bais, H. P. (2011). The role of ABC transporters in kin recognition in *Arabidopsis thaliana*. *Plant Signal. Behav.* **6**, 1154-1161. doi:10.4161/psb.6.8.15907
- Bombail, V. (2019). Perception and emotions: on the relationships between stress and olfaction. *Appl. Anim. Behav. Sci.* **212**, 98-108. doi:10.1016/j.applanim.2018.12.013
- Bombail, V., Barret, B., Raynaud, A., Jérôme, N., Saint-Albin, A., Ridder, C., Collin, A., Leterrier, C., Guilloteau, L. A. and Nielsen, B. L. (2018). In search of stress odours across species: behavioural responses of rats to faeces from chickens and rats subjected to various types of stressful events. *Appl. Anim. Behav. Sci.* **205**, 216-226. doi:10.1016/j.applanim.2017.10.013
- Borghi, L., Kang, J., Ko, D., Lee, Y. and Martinoia, E. (2015). The role of ABCG-type ABC transporters in phytohormone transport. *Biochem. Soc. Trans.* **43**, 924-930. doi:10.1042/BST20150106
- Brandl, H. B., Pruessner, J. C. and Farine, D. R. (2022). The social transmission of stress in animal collectives. *Proc. Biol. Sci.* **289**, 20212158.
- Brechbühl, J., Moine, F., Klaey, M., Nenniger-Tosato, M., Hurni, N., Sporkert, F., Giroud, C. and Broillet, M.-C. (2013). Mouse alarm pheromone shares structural similarity with predator scents. *Proc. Natl. Acad. Sci. USA* **110**, 4762-4767. doi:10.1073/pnas.1214249110
- Brown, G. E. (2003). Learning about danger: chemical alarm cues and local risk assessment in prey fishes. *Fish Fish. (Oxf)* **4**, 227-234. doi:10.1046/j.1467-2979.2003.00132.x
- Brown, G. E. and Smith, R. J. F. (1997). Conspecific skin extracts elicit antipredator responses in juvenile rainbow trout (*Oncorhynchus mykiss*). *Can. J. Zool.* **75**, 1916-1922. doi:10.1139/z97-821
- Brown, G. E., Poirier, J.-F. and Adrian, J. C., Jr (2004). Assessment of local predation risk: the role of subthreshold concentrations of chemical alarm cues. *Behav. Ecol.* **15**, 810-815. doi:10.1093/beheco/arh084
- Brunetti, A. E., Carnevale Neto, F., Vera, M. C., Taboada, C., Pavarini, D. P., Bauermeister, A. and Lopes, N. P. (2018). An integrative omics perspective for the analysis of chemical signals in ecological interactions. *Chem. Soc. Rev.* **47**, 1574-1591. doi:10.1039/C7CS00368D
- Burger, F. A., Terhaar, J. and Fröllicher, T. L. (2022). Compound marine heatwaves and ocean acidity extremes. *Nat. Commun.* **13**, 4722. doi:10.1038/s41467-022-32120-7
- Burks, R. L. and Lodge, D. M. (2002). Cued in: advances and opportunities in freshwater chemical ecology. *J. Chem. Ecol.* **28**, 1901-1917. doi:10.1023/A:1020785525081
- Cao, X. and Li, W. (2020). Embryonic substances induce alarm response in adult zebrafish (*Danio rerio*). *J. Fish Biol.* **97**, 225-230. doi:10.1111/jfb.14354
- Carnevali, L., Montano, N., Tobaldini, E., Thayer, J. F. and Sgoifo, A. (2020). The contagion of social defeat stress: Insights from rodent studies. *Neurosci. Biobehav. Rev.* **111**, 12-18. doi:10.1016/j.neubiorev.2020.01.011
- Chivers, D. P. and Smith, R. J. F. (1998). Chemical alarm signalling in aquatic predator-prey systems: a review and prospectus. *Écoscience* **5**, 338-352. doi:10.1080/11956860.1998.11682471
- Chivers, D. P., Brown, G. E. and Ferrari, M. C. O. (2012). The evolution of alarm substances and disturbance cues in aquatic animals. In *Chemical Ecology in Aquatic Systems* (ed. C. Brönmark and L.-A. Hansson), pp. 127-139. Oxford University Press.
- Coombs, S., Bleckmann, H., Fay, R. R. and Popper, A. N. (eds) (2013). *The Lateral Line System*, 2014 edn. New York: Springer.
- Cooper, G. M. (2000). Transport of small molecules. In *The Cell: A Molecular Approach*, 2nd edn. Sinauer Associates.
- Corey, D. P., García-Añoveros, J., Holt, J. R., Kwan, K. Y., Lin, S.-Y., Vollrath, M. A., Amalfitano, A., Cheung, E. L.-M., Derfler, B. H., Duggan, A. et al. (2004). TRPA1 is a candidate for the mechanosensitive transduction channel of vertebrate hair cells. *Nature* **432**, 723-730. doi:10.1038/nature03066
- Cowan, Z.-L., Green, L., Clark, T. D., Blewett, T. A., De Bonville, J., Gagnon, T., Hoots, E., Kuchenmüller, L., Leeuwis, R. H. J., Navajas Acedo, J. et al. (2024). Global change and premature hatching of aquatic embryos. *Glob. Chang. Biol.* **30**, e17488. doi:10.1111/gcb.17488
- Crane, A. L., Bairos-Novak, K. R., Goldman, J. A. and Brown, G. E. (2022). Chemical disturbance cues in aquatic systems: a review and prospectus. *Ecol. Monogr.* **92**, e01487. doi:10.1002/ecm.1487
- Cui, J., Ding, Y., Chen, S., Zhu, X., Wu, Y., Zhang, M., Zhao, Y., Li, T.-R. R., Sun, L. V., Zhao, S. et al. (2016). Disruption of Gpr45 causes reduced hypothalamic POMC expression and obesity. *J. Clin. Invest.* **126**, 3192-3206. doi:10.1172/JCI85676
- Dall, S. R. X., Giraldeau, L.-A., Olsson, O., McNamara, J. M. and Stephens, D. W. (2005). Information and its use by animals in evolutionary ecology. *Trends Ecol. Evol.* **20**, 187-193. doi:10.1016/j.tree.2005.01.010
- Darwin, C. (1871). *The Descent of Man and Selection in Relation to Sex*. London: John Murray.
- de Groot, J. H. B. and Smeets, M. A. M. (2017). Human fear chemosignaling: evidence from a meta-analysis. *Chem. Senses* **42**, 663-673. doi:10.1093/chemse/bjx049
- Deis, R., Lerer-Goldshtein, T., Baiko, O., Eyal, Z., Brenman-Begin, D., Goldsmith, M., Kaufmann, S., Heinig, U., Dong, Y., Lushchekina, S. et al. (2025). Genetic control over biogenic crystal morphogenesis in zebrafish. *Nat. Chem. Biol.* **21**, 383-392. doi:10.1038/s41589-024-01722-1
- Desban, L., Roussel, J., Mirat, O., Lejeune, F.-X., Keiser, L., Michalski, N. A. and Wyart, C. (2022). Lateral line hair cells integrate mechanical and chemical cues to orient navigation. *bioRxiv*. doi:10.1101/2022.08.31.505989
- Dijkgraaf, S. (1963). The functioning and significance of the lateral-line organs. *Biol. Rev. Camb. Philos. Soc.* **38**, 51-105. doi:10.1111/j.1469-185X.1963.tb00654.x
- Doney, S. C., Fabry, V. J., Feely, R. A. and Kleypas, J. A. (2009). Ocean acidification: the other CO₂ problem. *Ann. Rev. Mar. Sci.* **1**, 169-192. doi:10.1146/annurev.marine.010908.163834

- Elkon, R., Milon, B., Morrison, L., Shah, M., Vijayakumar, S., Racherla, M., Leitch, C. C., Silipino, L., Hadi, S., Weiss-Gayet, M. et al. (2015). RFX transcription factors are essential for hearing in mice. *Nat. Commun.* **6**, 8549. doi:10.1038/ncomms9549
- Fadok, V. A., Bratton, D. L., Frasch, S. C., Warner, M. L. and Henson, P. M. (1998). The role of phosphatidylserine in recognition of apoptotic cells by phagocytes. *Cell Death Differ.* **5**, 551-562. doi:10.1038/sj.cdd.4400404
- Federle, M. J. and Bassler, B. L. (2003). Interspecies communication in bacteria. *J. Clin. Invest* **112**, 1291-1299. doi:10.1172/JCI20195
- Ferrari, M. C. O., Wisenden, B. D. and Chivers, D. P. (2010). Chemical ecology of predator-prey interactions in aquatic ecosystems: a review and prospectus. *Can. J. Zool.* **88**, 698-724. doi:10.1139/Z10-029
- Feugere, L., Angell, L., Fagents, J., Nightingale, R., Rowland, K., Skinner, S., Hardege, J., Bartels-Hardege, H. and Wollenberg Valero, K. C. (2021a). Behavioural stress propagation in benthic invertebrates caused by acute pH drop-induced metabolites. *Front. Mar. Sci.* **8**, 773870. doi:10.3389/fmars.2021.773870
- Feugere, L., Scott, V. F., Rodriguez-Barucg, Q., Beltran-Alvarez, P. and Wollenberg Valero, K. C. (2021b). Thermal stress induces a positive phenotypic and molecular feedback loop in zebrafish embryos. *J. Therm. Biol.* **102**, 103114. doi:10.1016/j.jtherbio.2021.103114
- Feugere, L., Bates, A., Emagbetere, T., Chapman, E., Malcolm, L. E., Bulmer, K., Hardege, J., Beltran-Alvarez, P. and Wollenberg Valero, K. C. (2023). Heat induces multiomic and phenotypic stress propagation in zebrafish embryos. *PNAS Nexus* **2**, gad137. doi:10.1093/pnasnexus/pgad137
- Feugere, L., Silva De Freitas, C., Bates, A., Storey, K. B., Beltran-Alvarez, P. and Wollenberg Valero, K. C. (2025a). Social context prevents heat hormetic effects against mutagens during fish development. *FEBS Lett.* **599**, 2107-2128. doi:10.1002/1873-3468.70047
- Feugere, L., Fernandes, J. F., Guscetti, E., Melo, T., Aveiro, S., Chabot, D., Calado, R., Domingues, R., Madeira, D. and Calosi, P. (2025b). Non-additive phospholipidomic responses to ocean warming and acidification drive intraspecific variation in cell membrane vulnerability in a marine ectotherm. *Environ. Res.* **283**, 122135. doi:10.1016/j.envres.2025.122135
- Fletcher, N., Terschak, J. A., Bartels-Hardege, H. D., Bublitz, R., Schirmacher, P. and Hardege, J. D. (2022). A pheromone bouquet controls the reproductive behaviour of the male shore crab, *Carcinus maenas*. *Aquat. Ecol.* **56**, 419-427. doi:10.1007/s10452-021-09930-w
- Fortes-Marco, L., Lanuza, E. and Martinez-Garcia, F. (2013). Of pheromones and kairomones: what receptors mediate innate emotional responses? Pheromones and kairomones. *Anat. Rec.* **296**, 1346-1363. doi:10.1002/ar.22745
- Frantz, C., Stewart, K. M. and Weaver, V. M. (2010). The extracellular matrix at a glance. *J. Cell Sci.* **123**, 4195-4200. doi:10.1242/jcs.023820
- Gabius, H.-J., Siebert, H.-C., André, S., Jiménez-Barbero, J. and Rüdiger, H. (2004). Chemical biology of the sugar code. *ChemBiochem* **5**, 740-764. doi:10.1002/cbic.200300753
- Gagliano, M., Renton, M., Duvdevani, N., Timmins, M. and Mancuso, S. (2012). Out of sight but not out of mind: alternative means of communication in plants. *PLoS ONE* **7**, e37382. doi:10.1371/journal.pone.0037382
- Geraldo, L. H. M., Spohr, T. C. L. d. S., do Amaral, R. F., da Fonseca, A. C. C., Garcia, C., Mendes, F. d. A., Freitas, C., dosSantos, M. F. and Lima, F. R. S. (2021). Role of lysophosphatidic acid and its receptors in health and disease: novel therapeutic strategies. *Signal Transduct. Target. Ther.* **6**, 45. doi:10.1038/s41392-020-00367-5
- Gilmour, K. M. and Bard, B. (2022). Social buffering of the stress response: insights from fishes. *Biol. Lett.* **18**, 20220332. doi:10.1098/rsbl.2022.0332
- Godinho, C. P., Costa, R. and Sá-Correia, I. (2021). The ABC transporter Pdr18 is required for yeast thermotolerance due to its role in ergosterol transport and plasma membrane properties. *Environ. Microbiol.* **23**, 69-80. doi:10.1111/1462-2920.15253
- Goldberg, J. K., Wallace, A. K. and Weiss, S. L. (2017). Skin lipids of the striped plateau lizard (*Sceloporus virgatus*) correlate with female receptivity and reproductive quality alongside visual ornaments. *Sci. Nat.* **104**, 81. doi:10.1007/s00114-017-1503-3
- Goldman, J. A., Singh, A., Demers, E. E. M., Feyten, L. E. A. and Brown, G. E. (2019). Does donor group size matter? The response of guppies (*Poecilia reticulata*) and convict cichlids (*Amatitlania nigrofasciata*) to disturbance cues from conspecific and heterospecific donors. *Can. J. Zool.* **97**, 319-325. doi:10.1139/cjz-2018-0170
- Goldman, J. A., Feyten, L. E. A., Ramnarine, I. W. and Brown, G. E. (2020). Sender and receiver experience alters the response of fish to disturbance cues. *Curr. Zool.* **66**, 255-261. doi:10.1093/cz/zoz050
- Grunst, A. S. and Grunst, M. L. (2024). Animal personality in multiple stressor environments: the evolutionary ecology of among-individual differences in responses to stressor suites. *Proc. Biol. Sci.* **291**, 20241620.
- Guscetti, E., Spicer, J. I. and Calosi, P. (2019). The importance of inter-individual variation in predicting species' responses to global change drivers. *Ecol. Evol.* **9**, 4327-4339. doi:10.1002/ece3.4810
- Halpern, B. S., Frazier, M., Potapenko, J., Casey, K. S., Koenig, K., Longo, C., Lowndes, J. S., Rockwood, R. C., Selig, E. R., Selkoe, K. A. et al. (2015). Spatial and temporal changes in cumulative human impacts on the world's ocean. *Nat. Commun.* **6**, 7615. doi:10.1038/ncomms8615
- Hamdani, E. H. and Døving, K. B. (2007). The functional organization of the fish olfactory system. *Prog. Neurobiol.* **82**, 80-86. doi:10.1016/j.pneurobio.2007.02.007
- Hamdoun, A. and Epel, D. (2007). Embryo stability and vulnerability in an always changing world. *Proc. Natl. Acad. Sci. USA* **104**, 1745-1750. doi:10.1073/pnas.0610108104
- Hashiguchi, Y. and Nishida, M. (2007). Evolution of trace amine associated receptor (TAAR) gene family in vertebrates: lineage-specific expansions and degradations of a second class of vertebrate chemosensory receptors expressed in the olfactory epithelium. *Mol. Biol. Evol.* **24**, 2099-2107. doi:10.1093/molbev/msm140
- Hay, M. E. (2009). Marine chemical ecology: chemical signals and cues structure marine populations, communities, and ecosystems. *Ann. Rev. Mar. Sci.* **1**, 193-212. doi:10.1146/annurev.marine.010908.163708
- Hazel, J. R. (1995). Thermal adaptations in biological membranes: is homeoviscous adaptation the explanation? *Annu. Rev. Physiol.* **57**, 19-42. doi:10.1146/annurev.ph.57.030195.000315
- Hazel, J. R. (1997). Thermal adaptation in biological membranes: beyond homeoviscous adaptation. In *Thermobiology* (ed. J. S. Willis), pp. 57-101. Elsevier.
- Hazlett, B. A. (1985). Disturbance pheromones in the crayfish *Orconectes virilis*. *J. Chem. Ecol.* **11**, 1695-1711. doi:10.1007/BF01012121
- Hebets, E. A., Barron, A. B., Balakrishnan, C. N., Hauber, M. E., Mason, P. H. and Hoke, K. L. (2016). A systems approach to animal communication. *Proc. Biol. Sci.* **283**, 20152889. doi:10.1098/rspb.2015.2889
- Hews, D. K. (1988). Alarm response in larval western toads, *Bufo boreas*: release of larval chemicals by a natural predator and its effect on predator capture efficiency. *Anim. Behav.* **36**, 125-133. doi:10.1016/S0003-3472(88)80255-0
- Holm, H. C., Fredricks, H. F., Bent, S. M., Lowenstein, D. P., Ossolinski, J. E., Becker, K. W., Johnson, W. M., Schrage, K. and Van Mooy, B. A. S. (2022). Global ocean lipidomes show a universal relationship between temperature and lipid unsaturation. *Science* **376**, 1487-1491. doi:10.1126/science.abn7455
- Intergovernmental Panel on Climate Change (IPCC) (2019). *Special Report on the Ocean and Cryosphere in a Changing Climate* (ed. H.-O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Nicolai, A. Okem and others). Cambridge: Cambridge University Press. doi:10.1017/9781009157964
- Jin, P., Liang, Z., Lu, H., Pan, J., Li, P., Huang, Q., Guo, Y., Zhong, J., Li, F., Wan, J. et al. (2021). Lipid remodeling reveals the adaptations of a marine diatom to ocean acidification. *Front. Microbiol.* **12**, 748445. doi:10.3389/fmicb.2021.748445
- Jordão, L. and Volpato, G. (2000). Chemical transfer of warning information in non-injured fish. *Behaviour* **137**, 681-690. doi:10.1163/156853900502286
- Jorissen, H., Galand, P. E., Bonnard, I., Meiling, S., Raviglione, D., Meistertzheim, A.-L., Hédouin, L., Banaigs, B., Payri, C. E. and Nugues, M. M. (2021). Coral larval settlement preferences linked to crustose coralline algae with distinct chemical and microbial signatures. *Sci. Rep.* **11**, 14610. doi:10.1038/s41598-021-94096-6
- Kamio, M., Yambe, H. and Fusetani, N. (2022). Chemical cues for intraspecific chemical communication and interspecific interactions in aquatic environments: applications for fisheries and aquaculture. *Fish. Sci.* **88**, 203-239. doi:10.1007/s12562-021-01563-0
- Kandel, E. R., Schwartz, J. H., Jessell, T. M., Siegelbaum, S. A. and Hudspeth, A. J. (2000). *Principles of Neural Science*, 5th edn (ed. E. R. Kandel, J. H. Schwartz, T. M. Jessell, S. A. Siegelbaum and A. J. Hudspeth), p. 165. New York: McGraw-Hill Medical.
- Karakach, T. K., Huenupui, E. C., Soo, E. C., Walter, J. A. and Afonso, L. O. B. (2009). ¹H-NMR and mass spectrometric characterization of the metabolic response of juvenile Atlantic salmon (*Salmo salar*) to long-term handling stress. *Metabolomics* **5**, 123-137. doi:10.1007/s11306-008-0144-0
- Kassmer, S. H., Rodriguez, D. and De Tomaso, A. W. (2020). Evidence that ABC transporter-mediated autocrine export of an eicosanoid signaling molecule enhances germ cell chemotaxis in the colonial tunicate *Botryllus schlosseri*. *Development* **147**, dev184663. doi:10.1242/dev.184663
- Kim, S.-H., Turnbull, J. and Guimond, S. (2011). Extracellular matrix and cell signalling: the dynamic cooperation of integrin, proteoglycan and growth factor receptor. *J. Endocrinol.* **209**, 139-151. doi:10.1530/JOE-10-0377
- Kirsten, K., Fior, D., Kreutz, L. C. and Barcellos, L. J. G. (2018). First description of behavior and immune system relationship in fish. *Sci. Rep.* **8**, 846. doi:10.1038/s41598-018-19276-3
- Kiss, K., Brozik, A., Kucsma, N., Toth, A., Gera, M., Berry, L., Vallentin, A., Vial, H., Vidal, M. and Szakacs, G. (2012). Shifting the paradigm: the putative mitochondrial protein ABCB6 resides in the lysosomes of cells and in the plasma membrane of erythrocytes. *PLoS ONE* **7**, e37378. doi:10.1371/journal.pone.0037378
- Konishi, M. (1970). Evolution of design features in the coding of species-specificity. *Am. Zool.* **10**, 67-72. doi:10.1093/icb/10.1.67
- Laidre, M. E. and Johnstone, R. A. (2013). Animal signals. *Curr. Biol.* **23**, R829-R833. doi:10.1016/j.cub.2013.07.070
- Landshützer, P., Gruber, N., Bakker, D. C. E., Stemmler, I. and Six, K. D. (2018). Strengthening seasonal marine CO₂ variations due to increasing atmospheric CO₂. *Nat. Clim. Chang.* **8**, 146-150. doi:10.1038/s41558-017-0057-x

- Lauritano, C. and Ianora, A. (2020). Chemical defense in marine organisms. *Mar. Drugs* **18**, 518. doi:10.3390/md18100518
- Lee, S. S., Park, J. G., Jang, E., Choi, S. H., Kim, S., Kim, J. W. and Jin, M. S. (2023). W546 stacking disruption traps the human porphyrin transporter ABCB6 in an outward-facing transient state. *Commun. Biol.* **6**, 960. doi:10.1038/s42003-023-05339-3
- Le Guellec, D., Morvan-Dubois, G. and Sire, J.-Y. (2004). Skin development in bony fish with particular emphasis on collagen deposition in the dermis of the zebrafish (*Danio rerio*). *Int. J. Dev. Biol.* **48**, 217–231. doi:10.1387/ijdb.15272388
- Li, Z. and Nair, S. K. (2012). Quorum sensing: how bacteria can coordinate activity and synchronize their response to external signals? *Protein Sci.* **21**, 1403–1417. doi:10.1002/pro.2132
- Li, Z. and Vance, D. E. (2008). Phosphatidylcholine and choline homeostasis. *J. Lipid Res.* **49**, 1187–1194. doi:10.1194/jlr.R700019-JLR200
- Li, J., Mack, J. A., Souren, M., Yaksi, E., Higashijima, S.-I., Mione, M., Fetcho, J. R. and Friedrich, R. W. (2005). Early development of functional spatial maps in the zebrafish olfactory bulb. *J. Neurosci.* **25**, 5784–5795. doi:10.1523/JNEUROSCI.0922-05.2005
- Li, K., Brant, C. O., Huertas, M., Hessler, E. J., Mezei, G., Scott, A. M., Hoye, T. R. and Li, W. (2018). Fatty-acid derivative acts as a sea lamprey migratory pheromone. *Proc. Natl. Acad. Sci. USA* **115**, 8603–8608. doi:10.1073/pnas.1803169115
- Li, Y., Yan, Z., Lin, A., Yang, X., Li, X., Yin, X., Li, W. and Li, K. (2024). Epidermal oysterols function as alarm substances in zebrafish. *iScience* **27**, 109660. doi:10.1016/j.isci.2024.109660
- Liberles, S. D. (2009). Trace amine-associated receptors are olfactory receptors in vertebrates. *Ann. N. Y. Acad. Sci.* **1170**, 168–172. doi:10.1111/j.1749-6632.2009.04014.x
- Liberles, S. D. and Buck, L. B. (2006). A second class of chemosensory receptors in the olfactory epithelium. *Nature* **442**, 645–650. doi:10.1038/nature05066
- Liu, C. and Kakeya, H. (2020). Cryptic chemical communication: secondary metabolic responses revealed by microbial co-culture. *Chem. Asian J.* **15**, 327–337. doi:10.1002/asia.201901505
- Lucon-Xiccato, T., Di Mauro, G., Bisazza, A. and Bertolucci, C. (2020). Alarm cue-mediated response and learning in zebrafish larvae. *Behav. Brain Res.* **380**, 112446. doi:10.1016/j.bbr.2019.112446
- Mangiacotti, M., Baeckens, S., Fumagalli, M., Martín, J., Scali, S. and Sacchi, R. (2023). Protein-lipid association in lizard chemical signals. *Integr. Org. Biol.* **5**, obad016. doi:10.1093/iob/obad016
- Marchese, A., Sawdzargo, M., Nguyen, T., Cheng, R., Heng, H. H., Nowak, T., Im, D. S., Lynch, K. R., George, S. R. and O'Dowd, B. F. (1999). Discovery of three novel orphan G-protein-coupled receptors. *Genomics* **56**, 12–21. doi:10.1006/geno.1998.5655
- Mathuru, A. S., Kibat, C., Cheong, W. F., Shui, G., Wenk, M. R., Friedrich, R. W. and Jesuthasan, S. (2012). Chondroitin fragments are odorants that trigger fear behavior in fish. *Curr. Biol.* **22**, 538–544. doi:10.1016/j.cub.2012.01.061
- Matsumura, K., Matsunaga, S. and Fusetani, N. (2007). Phosphatidylcholine profile-mediated group recognition in catfish. *J. Exp. Biol.* **210**, 1992–1999. doi:10.1242/jeb.02777
- McDougall, T. J. and Barker, P. M. (2011). Getting started with TEOS-10 and the Gibbs Seawater (GSW) Oceanographic Toolbox. SCOR/IAPSO WG127. TEOS-10.
- McLean, S., Nichols, D. S. and Davies, N. W. (2021). Volatile scent chemicals in the urine of the red fox, *Vulpes vulpes*. *PLoS ONE* **16**, e0248961. doi:10.1371/journal.pone.0248961
- Melanson, C. A., Lamarre, S. G. and Currie, S. (2023). Social experience influences thermal sensitivity: lessons from an amphibious mangrove fish. *J. Exp. Biol.* **226**, jeb245656. doi:10.1242/jeb.245656
- Miyares, R. L., de Rezende, V. B. and Farber, S. A. (2014). Zebrafish yolk lipid processing: a tractable tool for the study of vertebrate lipid transport and metabolism. *Dis. Model. Mech.* **7**, 915–927. doi:10.1242/dmm.015800
- Miyazaki, M., Yamashita, T., Suzuki, Y., Saito, Y., Soeta, S., Taira, H. and Suzuki, A. (2006). A major urinary protein of the domestic cat regulates the production of feline, a putative pheromone precursor. *Chem. Biol.* **13**, 1071–1079. doi:10.1016/j.chembiol.2006.08.013
- Moros, M., Fergola, E., Marchesano, V., Mutarelli, M., Tommasini, G., Miedziak, B., Palumbo, G., Ambrosone, A., Tino, A. and Tortiglione, C. (2021). The aquatic invertebrate *Hydra vulgaris* releases molecular messages through extracellular vesicles. *Front. Cell Dev. Biol.* **9**, 788117. doi:10.3389/fcell.2021.788117
- Muñoz, D. J., Thorne, P. R. and Housley, G. D. (1999). P2X receptor-mediated changes in cochlear potentials arising from exogenous adenosine 5'-triphosphate in endolymph. *Hear. Res.* **138**, 56–64. doi:10.1016/S0378-5955(99)00151-3
- Nagahara, N., Nagano, M., Ito, T., Shimamura, K., Akimoto, T. and Suzuki, H. (2013). Antioxidant enzyme, 3-mercaptopyruvate sulfurtransferase-knockout mice exhibit increased anxiety-like behaviors: a model for human mercaptolactate-cysteine disulfiduria. *Sci. Rep.* **3**, 1986. doi:10.1038/srep01986
- Nie, L., Cai, S.-Y., Shao, J.-Z. and Chen, J. (2018). Toll-like receptors, associated biological roles, and signaling networks in non-mammals. *Front. Immunol.* **9**, 1523. doi:10.3389/fimmu.2018.01523
- Niu, L., Li, H., Song, Z., Dong, B., Cao, H., Liu, T., Du, T., Yang, W., Amin, R., Wang, L. et al. (2021). The functional analysis of ABCG transporters in the adaptation of pigeon pea (*Cajanus cajan*) to abiotic stresses. *PeerJ* **9**, e10688. doi:10.7717/peerj.10688
- Nozawa, Y. (2011). Adaptive regulation of membrane lipids and fluidity during thermal acclimation in *Tetrahymena*. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* **87**, 450–462. doi:10.2183/pjab.87.450
- Ohnstad, H., Jones, A. M., Howard, B., Schirmmacher, P., Bartels-Hardege, H. D. and Hardege, J. D. (2024). Effects of pH on olfactory behaviours in male shore crabs, *Carcinus maenas*. *Animals (Basel)* **14**, 948. doi:10.3390/ani14060948
- Parra, K. V., Adrian, J. C., Jr and Gerlai, R. (2009). The synthetic substance hypoxanthine 3-N-oxide elicits alarm reactions in zebrafish (*Danio rerio*). *Behav. Brain Res.* **205**, 336–341. doi:10.1016/j.bbr.2009.06.037
- Paulsson, M. and Wagener, R. (2018). Matrilins. *Methods Cell Biol.* **143**, 429–446. doi:10.1016/bs.mcb.2017.08.018
- Pelka, K. E., Henn, K., Keck, A., Sapel, B. and Braunbeck, T. (2017). Size does matter – determination of the critical molecular size for the uptake of chemicals across the chorion of zebrafish (*Danio rerio*) embryos. *Aquat. Toxicol.* **185**, 1–10. doi:10.1016/j.aquatox.2016.12.015
- Pfister, P. and Rodriguez, I. (2005). Olfactory expression of a single and highly variable V1r pheromone receptor-like gene in fish species. *Proc. Natl. Acad. Sci. USA* **102**, 5489–5494. doi:10.1073/pnas.0402581102
- Pichot, R., Watson, R. L. and Norton, I. T. (2013). Phospholipids at the interface: current trends and challenges. *Int. J. Mol. Sci.* **14**, 11767–11794. doi:10.3390/ijms140611767
- Poulin, R. X., Lavoie, S., Siegel, K., Gaul, D. A., Weissburg, M. J. and Kubanek, J. (2018). Chemical encoding of risk perception and predator detection among estuarine invertebrates. *Proc. Natl. Acad. Sci. USA* **115**, 662–667. doi:10.1073/pnas.1713901115
- Pozzer, A. C., Gómez, P. A. and Weiss, J. (2022). Volatile organic compounds in aquatic ecosystems - Detection, origin, significance and applications. *Sci. Total Environ.* **838**, 156155. doi:10.1016/j.scitotenv.2022.156155
- Quick, K., Zhao, J., Eijkelkamp, N., Linley, J. E., Rugiero, F., Cox, J. J., Raouf, R., Gringhuis, M., Sexton, J. E., Abramowitz, J. et al. (2012). TRPC3 and TRPC6 are essential for normal mechanotransduction in subsets of sensory neurons and cochlear hair cells. *Open Biol.* **2**, 120068. doi:10.1098/rsob.120068
- Rakvács, Z., Kucsma, N., Gera, M., Igriczi, B., Kiss, K., Barna, J., Kovács, D., Vellai, T., Bencs, L., Reisecker, J. M., Szoboszlai, N. and Szakács, G. (2019). The human ABCB6 protein is the functional homologue of HMT-1 proteins mediating cadmium detoxification. *Cell. Mol. Life Sci.* **76**, 4131–4144.
- Regnier, F. E. (1971). Semiochemicals—structure and function. *Biol. Reprod.* **4**, 309–326. doi:10.1093/biolreprod/4.3.309
- Roggatz, C. C., Fletcher, N., Benoit, D. M., Algar, A. C., Doroff, A., Wright, B., Wollenberg Valero, K. C. and Hardege, J. D. (2019). Saxitoxin and tetrodotoxin bioavailability increases in future oceans. *Nat. Clim. Chang* **9**, 840–844. doi:10.1038/s41558-019-0589-3
- Roggatz, C. C., Saha, M., Blanchard, S., Schirmmacher, P., Fink, P., Verheggen, F. and Hardege, J. D. (2022). Becoming nose-blind: climate change impacts on chemical communication. *Glob. Chang. Biol.* **28**, 4495–4505. doi:10.1111/gcb.16209
- Russo, M. V. and McGavern, D. B. (2015). Immune surveillance of the CNS following infection and injury. *Trends Immunol.* **36**, 637–650. doi:10.1016/j.it.2015.08.002
- Sanogo, Y. O., Hankison, S., Band, M., Obregon, A. and Bell, A. M. (2011). Brain transcriptomic response of threespine sticklebacks to cues of a predator. *Brain Behav. Evol.* **77**, 270–285. doi:10.1159/000328221
- Sasse, J., Simon, S., Gübeli, C., Liu, G.-W., Cheng, X., Friml, J., Bouwmeester, H., Martinioia, E. and Borghi, L. (2015). Asymmetric localizations of the ABC transporter PaPDR1 trace paths of directional strigolactone transport. *Curr. Biol.* **25**, 647–655. doi:10.1016/j.cub.2015.01.015
- Sbarbati, A. and Osculati, F. (2006). Allelochemical communication in vertebrates: kairomones, allomones and synomones. *Cells Tissues Organs* **183**, 206–219. doi:10.1159/000096511
- Selander, E., Kubanek, J., Hamberg, M., Andersson, M. X., Cervin, G. and Pavia, H. (2015). Predator lipids induce paralytic shellfish toxins in bloom-forming algae. *Proc. Natl. Acad. Sci. USA* **112**, 6395–6400. doi:10.1073/pnas.1420154112
- Sinensky, M. (1974). Homeoviscous adaptation—a homeostatic process that regulates the viscosity of membrane lipids in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **71**, 522–525. doi:10.1073/pnas.71.2.522
- Smith, R. J. F. (1979). Alarm reaction of Iowa and Johnny darters (*Etheostoma*, Percidae, Pisces) to chemicals from injured conspecifics. *Can. J. Zool.* **57**, 1278–1282. doi:10.1139/z79-164
- Snyder, N. and Snyder, H. (1970). Alarm response of *Diadema antillarum*. *Science* **168**, 276–278. doi:10.1126/science.168.3928.276
- Solomon, D. J. (1977). A review of chemical communication in freshwater fish. *J. Fish Biol.* **11**, 363–376. doi:10.1111/j.1095-8649.1977.tb04130.x
- Song, M., Zhao, J., Wen, H.-S., Li, Y., Li, J.-F., Li, L.-M. and Tao, Y.-X. (2019). The impact of acute thermal stress on the metabolome of the black rockfish (*Sebastes schlegelii*). *PLoS ONE* **14**, e0217133. doi:10.1371/journal.pone.0217133
- Sörbo, B. (1987). 3-Mercaptopyruvate, 3-mercaptoplactate and mercaptoacetate. *Methods Enzymol.* **143**, 178–182. doi:10.1016/0076-6879(87)43033-4

- Souza, A. R. C., Kozłowski, E. O., Cerqueira, V. R., Castelo-Branco, M. T. L., Costa, M. L. and Pavão, M. S. G. (2007). Chondroitin sulfate and keratan sulfate are the major glycosaminoglycans present in the adult zebrafish *Danio rerio* (Chordata-Cyprinidae). *Glycoconj. J.* **24**, 521-530. doi:10.1007/s10719-007-9046-z
- Stensmyr, M. C. and Maderspacher, F. (2012). Pheromones: fish fear factor. *Curr. Biol.* **22**, R183-R186. doi:10.1016/j.cub.2012.02.025
- Stincon, A., Prigione, A., Cramer, T., Wamelink, M. M. C., Campbell, K., Cheung, E., Olin-Sandoval, V., Grüning, N.-M., Krüger, A., Tauqeer Alam, M. et al. (2015). The return of metabolism: biochemistry and physiology of the pentose phosphate pathway. *Biol. Rev. Camb. Philos. Soc.* **90**, 927-963. doi:10.1111/brv.12140
- Syed, Z. A., Zhang, L. and Ten Hagen, K. G. (2022). In vivo models of mucin biosynthesis and function. *Adv. Drug Deliv. Rev.* **184**, 114182. doi:10.1016/j.addr.2022.114182
- Taga, M. E. and Bassler, B. L. (2003). Chemical communication among bacteria. *Proc. Natl. Acad. Sci. USA* **100** Suppl. 2, 14549-14554. doi:10.1073/pnas.1934514100
- Takahashi, F. T. (1974). Chemical communication and chemoreception in the Dungeness crab, *Cancer magister*, and the lined shore crab, *Pachygrapsus crassipes*. PhD thesis, Oregon State University.
- Tang, D., Kang, R., Coyne, C. B., Zeh, H. J. and Lotze, M. T. (2012). PAMPs and DAMPs: signals that spur autophagy and immunity. *Immunol. Rev.* **249**, 158-175. doi:10.1111/j.1600-065X.2012.01146.x
- Telser, A. (2002). *Molecular Biology of the Cell*, Vol. 18, 4th edn. Shock.
- Theodoulou, F. L. and Kerr, I. D. (2015). ABC transporter research: going strong 40 years on. *Biochem. Soc. Trans.* **43**, 1033-1040. doi:10.1042/BST20150139
- Tikhonov, D. B. and Zhorov, B. S. (2018). Predicting structural details of the sodium channel pore basing on animal toxin studies. *Front. Pharmacol.* **9**, 880. doi:10.3389/fphar.2018.00880
- Török, Z., Crul, T., Maresca, B., Schütz, G. J., Viana, F., Dindia, L., Piotto, S., Brameshuber, M., Balogh, G., Péter, M. et al. (2014). Plasma membranes as heat stress sensors: from lipid-controlled molecular switches to therapeutic applications. *Biochim. Biophys. Acta* **1838**, 1594-1618. doi:10.1016/j.bbamem.2013.12.015
- Turner, C. R., Spike, M. and Magrath, R. D. (2023). The evolution of eavesdropping on heterospecific alarm calls: Relevance, reliability, and personal information. *Ecol. Evol.* **13**, e10272. doi:10.1002/ece3.10272
- van der Vlist, E. J., Arkesteijn, G. J. A., van de Lest, C. H. A., Stoorvogel, W., Nolte-’t Hoen, E. N. M. and Wauben, M. H. M. (2012). CD4⁺ T cell activation promotes the differential release of distinct populations of nanosized vesicles. *J. Extracell. Vesicles* **1**, 18364. doi:10.3402/jev.v1i0.18364
- Van Theemsche, K. M., Van de Sande, D. V., Snyders, D. J. and Labro, A. J. (2020). Hydrophobic drug/toxin binding sites in voltage-dependent K⁺ and Na⁺ channels. *Front. Pharmacol.* **11**, 735. doi:10.3389/fphar.2020.00735
- Varki, A. (2011). Letter to the Glyco-Forum: since there are PAMPs and DAMPs, there must be SAMPs? Glycan 'self-associated molecular patterns' dampen innate immunity, but pathogens can mimic them. *Glycobiology* **21**, 1121-1124. doi:10.1093/glycob/cwr087
- Verheggen, F. J., Haubruge, E. and Mescher, M. C. (2010). Alarm pheromones—chemical signaling in response to danger. In *Vitamins & Hormones* (ed. G. Litwack), pp. 215-239. Academic Press.
- von Frisch, K. (1938). Zur Psychologie des Fisch-Schwarmes. *Sci. Nat.* **26**, 601-606. doi:10.1007/BF01590598
- von Frisch, K. (1942). Ueber einen Schreckstoff der Fischhaut und seine biologische Bedeutung. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **29**, 46-145. doi:10.1007/BF00304445
- Vigh, L., Maresca, B. and Harwood, J. L. (1998). Does the membrane's physical state control the expression of heat shock and other genes? *Trends Biochem. Sci.* **23**, 369-374. doi:10.1016/S0968-0004(98)01279-1
- Webb, J. F. (2013). Morphological diversity, development, and evolution of the mechanosensory lateral line system. In *Springer Handbook of Auditory Research* (ed. A. B. Coffin and J. Sisneros), pp. 17-72. New York: Springer New York.
- Wisenden, B. D., Vollbrecht, K. A. and Brown, J. L. (2004). Is there a fish alarm cue? Affirming evidence from a wild study. *Anim. Behav.* **67**, 59-67. doi:10.1016/j.anbehav.2003.02.010
- Wyatt, T. D. (2014). *Pheromones and Animal Behavior: Chemical Signals and Signatures*. Cambridge University Press.
- Xiang, Z., Bo, X. and Burnstock, G. (1999). P2X receptor immunoreactivity in the rat cochlea, vestibular ganglion and cochlear nucleus. *Hear. Res.* **128**, 190-196. doi:10.1016/S0378-5955(98)00208-1
- Ye, Z., Lu, Y. and Wu, T. (2020). The impact of ATP-binding cassette transporters on metabolic diseases. *Nutr. Metab.* **17**, 61.
- Zhang, D., Terschak, J. A., Harley, M. A., Lin, J. and Hardege, J. D. (2011). Simultaneously hermaphroditic shrimp use lipophilic cuticular hydrocarbons as contact sex pheromones. *PLoS ONE* **6**, e17720. doi:10.1371/journal.pone.0017720
- Zhang, W., Xu, X., Li, J. and Shen, Y. (2022). Transcriptomic analysis of the liver and brain in grass carp (*Ctenopharyngodon idella*) under heat stress. *Mar. Biotechnol.* **24**, 856-870. doi:10.1007/s10126-022-10148-6